

Mosquito Management and Risk by Robert K D Peterson

In his inauguration speech on January 21, 2009, President Obama said "We'll restore science to its rightful place..." Although he was referring primarily to the role of science in economic and technology development, it is appropriate to extend his statement to include science's role in societal decisions about technology and the regulation of technology. In particular, what does restoring science to its rightful place mean when it comes to the regulation of technology? Because technology is based on science, scientific evidence must be afforded its proper place in decisions about the proper use of technology. Of course, in our democratic system, science does not – nor should not – serve as the sole arbiter of societal decisions about technologies and how to use them. These decisions should use science as a foundation, but then also incorporate economic, legal, ethical, aesthetic, and cultural factors.

What does "restoring science to its rightful place" have to do with mosquito management and risk? Quite a lot, actually. Since the infectious pathogen West Nile virus invaded the United States in 1999, causing the largest encephalitis disease epidemic in US history (1), renewed public attention has been focused on mosquito management. Most of this attention and concern has involved using insecticides as an outdoor space application targeting adult mosquitoes, often called "adulticiding" (2). The concerns have revolved around two major areas: the effectiveness of adulticiding, and the risks posed

by adulticiding. These concerns have led directly to opposition to adulticiding by organized activist groups and have led to misinformation and opinions that are not consistent with facts. The risks and benefits of adulticiding and other management tactics have been extensively studied. So, let's look at each of the two major areas identified above and see what the prevailing science has to say.

We have long known that using adulticides in outdoor space applications reduces populations of adult mosquitoes, although the results can be quite variable. What has been more uncertain is the effect of adulticiding on reducing pathogen infection rates in mosquitoes and on reducing disease incidence in people and other animals. Although it is reasonable to assume that if adulticiding reduces adult mosquito populations there will also be reductions in pathogen-infected mosquitoes and disease in people and other animals, research to evaluate this has been lacking until very recently. Results from these studies suggest that adulticiding has a significant effect on reducing pathogen infections in mosquitoes and disease (3-5).

What about the ecological and human-health risks from adulticiding? Before we can examine these issues, we need to take a step back and look more broadly at the scientific discipline of risk assessment. To assess risk, one needs to understand both *effect* and *exposure*. Risk is really nothing more than an interaction of these two factors (6). And, risk assessment is

simply the objective evaluation of risk in which the assumptions and uncertainties that are part of the assessment are clearly considered and presented (6,7).

In the case of adulticides, we need to know the toxicity of, and exposure to, the insecticide in question to properly estimate the risks. Like all chemical risk assessments, researchers have estimated or measured the exposure of a person or other organism to the insecticide and then compared that exposure to a threshold exposure level. The threshold level is usually determined by a regulatory agency such as the US Environmental Protection Agency (EPA). Threshold exposure levels usually incorporate safety factors that increase the protection of people and wildlife. In the case of exposure and risk to people, the threshold level is the exposure to the insecticide that has been shown in a series of studies with laboratory animals to have no toxic effects on individuals. For exposure and risk to non-target organisms, like wildlife, the threshold level varies and may be an exposure in which no toxic effects are observed, or it may be an exposure that is a small fraction of the concentration needed to kill 50% of the test-animal population. In other words, there may be a toxic effect, but it has been determined that it will not affect the populations of organisms in the short or long term.

What do the risk assessments that have been conducted for outdoor space applications have to say? Because risk assessment of insecticides is

dependent on the amount and frequency of exposure, the application rate tells us a lot about the resulting risk. Insecticides used for outdoor space applications are applied at a very low rate compared to agricultural and residential applications. For example, in agricultural applications permethrin is often applied at a rate thirty-five times greater than is applied for adult mosquito management. In a sense, then, the risk from outdoor space applications for adult mosquitoes is just a small fraction of the risk from agricultural applications. But is that risk acceptable or unacceptable?

The weight of scientific evidence from independent research and regulatory agency assessments strongly suggests that exposures to people from outdoor space applications of mosquito adulticides are well below threshold levels of concern (2,5,8-20). Epidemiology and biomonitoring studies (21-25) support the results from these risk assessments. Exposures to mammals, birds, fish, and aquatic invertebrates are also below levels of concern (19,26-35). These assessments include short- and long-term exposures from single and multiple applications. Risks are low even for many terrestrial insects because the applications target flying mosquitoes at night (26,27,32,35,36). Risks to people, mammals, and birds are extremely low. For example, a person's exposure to permethrin as a result of an outdoor space application would be less than one ten-thousandth of the threshold exposure level (8,19,37); see Figure 1.

What about the risks from other mosquito management tactics such as larvicides, personal repellents, and insecticide-treated clothing and bednets? Here,




Figure 1: Estimated exposure of a person to the adulticide permethrin (0.000025 mg/kg body weight), compared to the US EPA's Acceptable Daily Exposure Level (0.25 mg/kg body weight) and the No-Effect Level (25 mg/kg body weight). The Acceptable Daily Exposure Level is the amount of chemical that an individual can be exposed to over a lifetime without experiencing any toxic effects. The No-Effect Level is the exposure at which no toxic effects have been observed in a series of laboratory animal experiments.

too, research over many years and from many researchers indicates that exposures are below levels of concern, provided that these tactics are properly used (18,29,38-44). However, the use of the western mosquitofish, a biological control organism, may present unacceptable risks to fish and aquatic invertebrates where the fish is not currently endemic (45).

Mosquito management is essentially part of a broader public and environmental health enterprise. In mosquito management, we use an Integrated Pest Management (IPM) approach. This involves identification of mosquito species and surveillance of their populations. When populations of larvae, pupae, adults, or pathogen-infected adults reach pre-established threshold levels,

actions may be taken to lower those populations below the thresholds. These actions may involve several tactics. When adult mosquito or disease thresholds have been breached, adulticides may be used. As we have seen above, the use of adulticides by health professionals can reduce infected mosquitoes and resultant disease. And, as we have seen, when these adulticides are used in outdoor space applications, the risks are below levels of concern.

In the US, West Nile virus and many other pathogens are alien, invasive organisms. The diseases these pathogens cause pose human health and economic problems for our society, and they pose environmental problems for ecosystem functioning and biodiversity. As part of the public and

environmental health enterprise, mosquito management professionals are charged with the well-being of their particular local areas. If we are to put science in its rightful place, science should inform our understanding of risk and the societal decisions that need to be made about mosquito and disease management. The weight of scientific evidence shows that mosquitoes and the pathogens they carry can cause appreciable risks to public and environmental health. When mosquito populations need to be managed, IPM tactics are used that have been shown to be effective and to result in risks that are below levels of concern. Science, then, is put in its proper place by providing the facts that are used to make the best decisions to protect our health and the environment.

For more information and a complete list of citations from this article, see the West Nile Virus, Mosquito Management, and Risk website at: <http://landresources.montana.edu/WNV/>.

SELECTED REFERENCES CITED

Huhn GD, Sejvar JJ, Montgomery SP, and Dworkin MS. 2003. West Nile virus in the United States: an update on an emerging infectious disease. *American Family Physician* 68:653-660.

Peterson RKD, Macedo PA, and Davis RS. 2006. A human-health risk assessment for West Nile virus and insecticides used in mosquito management. *Environmental Health Perspectives* 114:366-372.

Elnaiem D-EA, Kelley K, Wright S, Laffey R, Yoshimura G, Reed M, Goodman G, Thiemann T, Reimer L, Reisen WK, and Brown D. 2008. Impact of aerial spraying of pyrethrin insecticide on *Culex pipiens*

and *Culex tarsalis* (Diptera: Culicidae) abundance and West Nile virus infection rates in an urban/suburban area of Sacramento County, California. *Journal of Medical Entomology* 45:751-757.

Carney RM, Husted S, Jean C, Glaser C, and Kramer V. 2008. Efficacy of aerial spraying of mosquito adulticide in reducing incidence of West Nile virus, California, 2005. *Emerging Infectious Diseases* 14:747-754.

Macedo PA, Schleier III JJ, Reed M, Kelley K, Goodman GW, Brown DA, and Peterson RKD. 2010. Evaluation of efficacy and human health risk of aerial ultra-low volume applications of pyrethrins and piperonyl butoxide for adult mosquito management in response to West Nile virus activity in Sacramento County, California. *Journal of the American Mosquito Control Association* 26:57-66.

NRC. 1983. Risk assessment in the federal government: managing the process. National Research Council, National Academy Press, Washington, DC.

NRC. 1996. Understanding risk: informing decisions in a democratic society. National Research Council, National Academy Press, Washington, DC.



Robert K D Peterson
Associate Professor
Department of Land Resources and Environmental Sciences
Montana State University
Bozeman, MT 59717
406-994-7927
bpeterson@montana.edu

FOUR STAR MICROBIAL BRIQUETS

—the silver bullet you've been waiting for.

FourStar Briquets are powerful ammunition for the sites you'd like to treat and forget about.

Three sizes available:

- 45 day briquets
- 90 day briquets
- 180 day briquets



To learn more about FourStar Briquets, contact ADAPCO toll free by phone at (800) 367-0659 by email at info@myadapco.com, or visit us at www.fourstarmicrobials.com.

Complete references for the article: **Peterson, R. K. D. 2010. Mosquito management and risk. *Wing Beats*, Fall 2010, 21(3):28-31.**

References Cited

1. Huhn GD, Sejvar JJ, Montgomery SP, and Dworkin MS (2003) West Nile virus in the United States: an update on an emerging infectious disease. *American Family Physician* 68:653-660.
2. Peterson RKD, Macedo PA, and Davis RS (2006) A human-health risk assessment for West Nile virus and insecticides used in mosquito management. *Environmental Health Perspectives* 114:366-372.
3. Elnaiem D-EA, Kelley K, Wright S, Laffey R, Yoshimura G, Reed M, Goodman G, Thiemann T, Reimer L, Reisen WK, and Brown D (2008) Impact of aerial spraying of pyrethrin insecticide on *Culex pipiens* and *Culex tarsalis* (Diptera: Culicidae) abundance and West Nile virus infection rates in an urban/suburban area of Sacramento County, California. *Journal of Medical Entomology* 45:751-757.
4. Carney RM, Husted S, Jean C, Glaser C, and Kramer V (2008) Efficacy of aerial spraying of mosquito adulticide in reducing incidence of West Nile virus, California, 2005. *Emerging Infectious Diseases* 14:747-754.
5. Macedo PA, Schleier III JJ, Reed M, Kelley K, Goodman GW, Brown DA, and Peterson RKD (2010) Evaluation of efficacy and human health risk of aerial ultra-low volume applications of pyrethrins and piperonyl butoxide for adult mosquito management in response to West Nile virus activity in Sacramento County, California. *Journal of the American Mosquito Control Association* 26:57-66.
6. NRC (1983) Risk assessment in the federal government: managing the process. National Research Council, National Academy Press, Washington, D.C.
7. NRC (1996) Understanding risk: informing decisions in a democratic society. National Research Council, National Academy Press, Washington, D.C.
8. Schleier III JJ, Macedo PA, Davis RS, Shama LM, and Peterson RKD (2009) A two-dimensional probabilistic acute human-health risk assessment of insecticide exposure after adult mosquito management. *Stochastic Environmental Research and Risk Assessment* 23:555-563.
9. Carr WC, Iyer P, and Gammon DW (2006) A dietary risk assessment of the pyrethroid insecticide resmethrin associated with its use for West Nile virus mosquito vector control in California. *The Scientific World Journal* 6:279-290.
10. NYCDOH. Adult mosquito control programs: environmental impact statement (EIS). New York City Department of Health: New York, NY, USA (2005).
11. Suffolk County. Draft generic environmental impact statement. Suffolk County Department of Public Works and Department of Health Services: Suffolk County, NY (2006).
12. USEPA. Reregistration eligibility decision (RED) for permethrin. U.S. Environmental Protection Agency: Washington D.C., pp. 1-95 (2006).
13. USEPA. Reregistration eligibility decision for resmethrin. U.S. Environmental Protection Agency: Washington D.C., USA (2006).
14. USEPA. Reregistration eligibility decision for pyrethrins. U.S. Environmental Protection Agency: Washington D.C., pp. 1-108 (2006).
15. USEPA. Reregistration eligibility decision for piperonyl butoxide (PBO). U.S. Environmental Protection Agency: Washington D.C., pp. 1-111 (2006).
16. USEPA. Reregistration eligibility decision for malathion. U.S. Environmental Protection Agency: Washington D.C., pp. 1-101 (2006).

17. USEPA. Interim reregistration eligibility decision for naled, Case No. 0092. U.S. Environmental Protection Agency: Washington D.C., pp. 1-130 (2006).
18. Macedo PA, Peterson RKD, and Davis RS (2007) Risk assessments for exposure of deployed military personnel to insecticides and personal protective measures used for disease-vector management. *Journal of Toxicology and Environmental Health, Part A* 70:1758–1771.
19. Schleier III JJ, and Peterson RKD (2010) Deposition and air concentrations of permethrin and naled used for adult mosquito management. *Archives of Environmental Contamination and Toxicology* 58:105-111.
20. Valcke M, Gosselin N, and Belleville D (2008) Human exposure to malathion during a possible vector-control intervention against West Nile Virus. II: evaluation of the toxicological risks using a probabilistic approach. *Human and Ecological Risk Assessment* 14:1138-1158.
21. Currier M, McNeill M, Campbell D, Newton N, Marr JS, Perry E, Berg SW, Barr DB, Luber GE, Kieszak SM, Rogers HS, Backer SC, Belson MG, Rubin C, Azziz-Baumgartner E, and Duprey ZH (2005) Human exposure to mosquito-control pesticides---Mississippi, North Carolina, and Virginia, 2002 and 2003. *MMWR Morbidity and Mortality Weekly Report* 54:529-532.
22. Kutz FW, and Strassman SC (1977) Human urinary metabolites of organophosphate insecticides following mosquito adulticiding. *Mosquito News* 37:211-218.
23. Karpati AM, Perrin MC, Matte T, Leighton J, Schwartz J, and Barr RG (2004) Pesticide spraying for West Nile virus control and emergency department asthma visits in New York City, 2000. *Environmental Health Perspectives* 112:1183-1187.
24. O'Sullivan BCY, Lafleur J, Fridal K, Hormozdi S, Schwartz S, Belt M, and Finkel M (2005) The effect of pesticide spraying on the rate and severity of ED asthma. *American Journal of Emergency Medicine* 23:463-467.
25. Duprey Z, Rivers S, Luber G, Becker A, Blackmore C, Barr D, Weerasekera G, Kieszak S, Flanders WD, and Rubin C (2008) Community aerial mosquito control and naled exposure. *Journal of the American Mosquito Control Association* 24:42-46.
26. Kwan JA, Novak MG, Hyles TS, and Niemela MK (2009) Mortality of nontarget arthropods from an aerial application of pyrethrins. *Journal of the American Mosquito Control Association* 25:218-220.
27. Antwi FB, and Peterson RKD (2009) Toxicity to non-target insects after exposure to δ -phenothrin and resmethrin. *Pest Management Science* 65:300-305.
28. Davis RS, Peterson RKD, and Macedo PA (2007) An ecological risk assessment for insecticides used in adult mosquito management. *Integrated Environmental Assessment and Management* 3:373-382.
29. Davis RS, and Peterson RKD (2008) Effects of single and multiple applications of mosquito insecticides on nontarget arthropods. *Journal of the American Mosquito Control Association* 24:270-280.
30. Schleier III JJ, Davis RS, Shama LM, Macedo PA, and Peterson RKD (2008) Equine risk assessment for insecticides used in adult mosquito management. *Human and Ecological Risk Assessment* 14:392-407.
31. Schleier III JJ, Peterson RKD, Macedo PA, and Brown DA (2008) Environmental concentrations, fate, and risk assessment of pyrethrins and piperonyl butoxide after aerial ultralow-volume applications for adult mosquito management. *Environmental Toxicology and Chemistry* 27:1063-1068.
32. Schleier III JJ, and Peterson RKD (2010) Toxicity and risk of permethrin and naled to non-target insects after adult mosquito management. *Ecotoxicology* 16:1140-1146.
33. Jensen T, Lawler SP, and Dritz DA (1999) Effects of ultra-low volume pyrethrin, malathion, and permethrin on nontarget invertebrates, sentinel mosquitoes, and mosquitofish in seasonally impounded wetlands. *Journal of the American Mosquito Control Association* 15:330-338.

34. Breaud TP, Farlow JE, Steelman CD, and Schilling PE (1977) Effects of the insect growth regulator methoprene on natural populations of aquatic organisms in Louisiana intermediate marsh habitats. *Mosquito News* 37:704-712.
35. Boyce WM, Lawler SP, Schultz JM, McCauley SJ, Kimsey LS, Niemela MK, Nielsen CF, and Reisen WK (2007) Nontarget effects of the mosquito adulticide pyrethrin applied aerially during a West Nile virus outbreak in an urban California environment. *Journal of the American Mosquito Control Association* 23:335-339.
36. Breidenbaugh MS, and de Szalay FA (2010) Effects of aerial applications of naled on nontarget insects at Parris Island, South Carolina. *Environmental Entomology* 39:591-599.
37. Schleier III JJ. Environmental concentrations, fate, and risk assessment of insecticides used for adult mosquito management. Montana State University, Bozeman, MT, (2008).
38. Hershey AE, Lima AR, Niemi GJ, and Regal RR (1998) Effects of *Bacillus thuringiensis israelensis* (Bti) and methoprene on nontarget macroinvertebrates in Minnesota wetlands. *Ecological Applications* 8:41-60.
39. Barlow SM, Sullivan FM, and Lines J (2001) Risk assessment of the use of deltamethrin on bednets for the prevention of malaria. *Food and Chemical Toxicology* 39:407-422.
40. Antwi F, Shama LM, and Peterson RKD (2008) Risk assessments for the insect repellents DEET and picaridin. *Regulatory Toxicology and Pharmacology* 51:31-36.
41. Zaim M, Aitio A, and Nakashima N (2000) Safety of pyrethroid-treated mosquito nets. *Medical and Veterinary Entomology* 14:1-5.
42. Charbonneau CS, Drobney RD, and Rabeni CF (1994) Effects of *Bacillus thuringiensis* var. *israelensis* on nontarget benthic organisms in a lentic habitat and factors affecting the efficacy of the larvicide. *Environmental Toxicology and Chemistry* 13:267-269.
43. Lawler SP, Jensen T, Dritz DA, and Wichterman G (1999) Field efficacy and nontarget effects of the mosquito larvicides temephos, methoprene, and Bti in Florida mangrove swamps. *Journal of the American Mosquito Control Association* 15:446-452.
44. Niemi GJ, Hershey AE, Shannon L, Hanowski JM, Lima A, Axler RP, and Regal RR (1999) Ecological effects of mosquito control on zooplankton, insects, and birds. *Environmental Toxicology and Chemistry* 18:549-559.
45. Schleier III JJ, Sing SE, and Peterson RKD (2008) Regional ecological risk assessment for the introduction of *Gambusia affinis* (western mosquitofish) into Montana watersheds. *Biological Invasions* 10:1277-1287.